

Patient Name:

Date/Time:

Screened by:

#### COVID-19 SEVERITY:

- **Mild:** Includes fever, cough, sore throat, malaise, headache, muscle pain, nausea, vomiting, diarrhea, loss of taste and smell but who do not have shortness of breath, dyspnea, or abnormal chest imaging
- **Moderate:** Shows evidence of lower respiratory disease during clinical assessment or imaging and who have an oxygen saturation (SpO<sub>2</sub>)  $\geq$ 94% on room air at sea level
- **Severe** (unlikely to be encountered in the home setting) Individuals who have SpO<sub>2</sub>  $<$ 94% on room air at sea level, a ratio of arterial partial pressure of oxygen to fraction of inspired oxygen (PaO<sub>2</sub>/FiO<sub>2</sub>)  $<$ 300 mm Hg, a respiratory rate  $>$ 30 breaths/min, or lung infiltrates  $>$ 50%.

#### MODERATE AND SEVERE IMMUNOCOMPROMISING CONDITIONS:

Moderate and severe immunocompromising conditions and treatments include but are not limited to the following. The degree of immunocompromise for the patient is determined by the treating provider, and preventive actions should be tailored to each patient and situation.

- Active treatment for solid tumor and hematologic malignancies
- Receipt of solid-organ transplant and taking immunosuppressive therapy
- Receipt of chimeric antigen receptor (CAR)-T-cell therapy or hematopoietic cell transplant (HCT) (within 2 years of transplantation or taking immunosuppressive therapy)
- Moderate or severe primary immunodeficiency (e.g., CVID, Selective Antibody Deficiency, Subclass Deficiency, SCID)
- Advanced HIV infection (people with HIV and CD4 cell counts less than 200/mm<sup>3</sup>, history of an AIDS-defining illness without immune reconstitution, or clinical manifestations of symptomatic HIV) or untreated HIV infection
- Active treatment with high-dose corticosteroids (i.e., 20 mg or more of prednisone or equivalent per day when administered for 2 or more weeks), alkylating agents, antimetabolites (e.g., methotrexate, immunosuppressive drugs used for transplants/other reason (e.g., azathioprine/Imuran<sup>®</sup>, cyclosporine, mycophenolate/CellCept<sup>®</sup>, tacrolimus), cancer chemotherapeutic agents classified as severely immunosuppressive, tumor necrosis factor (TNF) blockers (e.g., infliximab/Inflectra<sup>®</sup>/Remicade<sup>®</sup>/Renflexis<sup>®</sup>, Enbrel<sup>®</sup>, Humira<sup>®</sup>, Cimzia<sup>®</sup>, and Simponi<sup>®</sup>), and other biologic agents that are immunosuppressive or immunomodulatory

Part 1. Patient Screen

Call patient no later than 24 hours before the visit.

Patient Name: \_\_\_\_\_

**QUESTION 1**

Is patient considered to be moderately / severely immunocompromised (see information box above)?      YES      NO

**QUESTION 2**

Have you tested positive for COVID-19 within the last 20 days?      YES (continue)      NO (proceed to Question 3)

- Positive test date: \_\_\_\_\_
- Date symptoms started: \_\_\_\_\_      [ ] No symptoms
- Is prescriber of Nufactor therapy aware?      YES      NO
- If severe symptoms, contact Nufactor

**QUESTION 2A**

If **NOT** moderately / severely immunocompromised

**Asymptomatic**

**Symptomatic**

- 10 days from positive test: \_\_\_\_\_  
(day 0 = specimen collection date; day 1 = the next full day)
- Advise patient to isolate from others and wear a well-fitting mask around others x 10 days after positive test.
- Advise patient to avoid places where unable to wear a mask.
- **If asymptomatic:**
  - End isolation/masking after 10 days.
  - May end isolation/masking earlier with testing if results are negative from at least two consecutive respiratory specimens collected 48 hours apart (total of two negative specimens) tested using an antigen test or Nucleic Acid Amplification Test (NAAT)/Polymerase Chain Reaction (PCR) test.
  - If symptoms develop, restart isolation at day 0 and follow symptomatic guidelines.

- **If symptomatic:**
  - End isolation/masking after 10 days if 1) At least 24 hours since last fever (without use of anti-pyretics) **and** 2) Symptoms have improved (excluding loss of taste and smell, which may persist for weeks or months after recovery).
  - May end isolation/masking earlier with testing if 1) At least 24 hours since last fever (without use of anti-pyretics) **and** 2) Symptoms have improved (excluding loss of taste and smell, which may persist for weeks or months after recovery) **and** 3) Results are negative from at least two consecutive respiratory specimens collected 48 hours apart (total of two negative specimens) tested using an antigen test or Nucleic Acid Amplification Test (NAAT)/Polymerase Chain Reaction (PCR) test.
  - If symptoms worsen or recur, restart isolation at day 0.
- Inform Nufactor
- If willing to make visit in the 10 day period when patient should isolate/mask, don full PPE (N-95 mask, eye protection/face shield, gloves, gown, and have patient don N-95 mask). Only the nurse and patient should be in the room during the visit.

Patient Name: \_\_\_\_\_

**QUESTION 2B**

**If moderately / severely immunocompromised      Asymptomatic      Symptomatic**

- 20 days from positive test: \_\_\_\_\_ (day 0 = specimen collection date; day 1 = the next full day)
- Advise patient to isolate from others and wear a well-fitting mask around others x 20 days after positive test.
- Advise patient to avoid places where unable to wear a mask.

**If asymptomatic/never developed symptoms:**

- End isolation/masking after 20 days.
- May end isolation/masking earlier with testing if results are negative from at least two consecutive respiratory specimens collected 48 hours apart (total of two negative specimens) tested using an antigen test or Nucleic Acid Amplification Test (NAAT)/Polymerase Chain Reaction (PCR) test.
- If symptoms develop, restart isolation at day 0 and follow symptomatic guidelines.

**If symptomatic:**

- End isolation/masking after 20 days if 1) At least 24 hours since last fever (without use of anti-pyretics) **and** 2) Symptoms have improved (excluding loss of taste and smell, which may persist for weeks or months after recovery).

- May end isolation/masking earlier with testing if 1) At least 24 hours since last fever (without use of anti-pyretics) **and** 2) Symptoms have improved (excluding loss of taste and smell, which may persist for weeks or months after recovery) **and** 3) Results are negative from at least two consecutive respiratory specimens collected 48 hours apart (total of two negative specimens) tested using an antigen test or Nucleic Acid Amplification Test (NAAT)/Polymerase Chain Reaction (PCR) test.

- If symptoms worsen or recur, restart isolation at day 0.
- Inform Nufactor
- If willing to make visit in the 20 day period when patient should isolate/mask, don full PPE (N-95 mask, eye protection/face shield, gloves, gown, and have patient don N-95 mask). Only the nurse and patient should be in the room during the visit.

**QUESTION 3**

**Do you have any COVID-19 symptoms that are not caused by another condition such as allergies?**

Contact Nufactor if unsure.    **YES** (continue)    **NO** (proceed to Question 4)

- Date symptoms started: \_\_\_\_\_
- If awaiting test result, date of test: \_\_\_\_\_
- Is prescriber of Nufactor therapy aware?    yes    no
- If severe symptoms, contact Nufactor.

**Check all that apply**

- |  |   |   |
|--|---|---|
| <input type="checkbox"/> Fever or chills                             | <input type="checkbox"/> Muscle or body aches       | <input type="checkbox"/> Congestion or runny nose |
| <input type="checkbox"/> Cough                                       | <input type="checkbox"/> Headache                   | <input type="checkbox"/> Nausea or vomiting       |
| <input type="checkbox"/> Shortness of breath or difficulty breathing | <input type="checkbox"/> New loss of taste or smell | <input type="checkbox"/> Diarrhea                 |
| <input type="checkbox"/> Fatigue                                     | <input type="checkbox"/> Sore throat                |   |

Patient Name:

**QUESTION 3A**

**If NOT moderately / severely immunocompromised**

- 10 days from symptom start date: \_\_\_\_\_ (day 0; day 1 = next full day)
- Advise patient to test as soon as possible.
- Advise patient to isolate from others and wear a well-fitting mask around others x 10 days since symptom onset (day 0 = symptom start date; day 1 = the next full day)
- Advise patient to avoid places where unable to wear a mask.
- End isolation/masking after 10 days if 1) At least 24 hours since last fever (without use of anti-pyretics) **and** 2) Symptoms have improved.
- May end isolation/masking earlier with testing if 1) At least 24 hours since last fever (without use of anti-pyretics) **and** 2) Symptoms have improved **and** 3) If using, a single negative Nucleic Acid Amplification Test (NAAT)/ Polymerase Chain Reaction (PCR) test is sufficient in most circumstances. If a higher level of clinical suspicion for COVID-19 infection exists, consider maintaining extending masking/isolation and confirming with a second negative NAAT/PCR test or if using an antigen test, a negative result should be confirmed by either a negative NAAT/PCR test or second negative antigen test taken 48 hours after the first negative test.
- Inform Nufactor
- If symptoms worsen or recur, restart isolation at day 0.
- If willing to make visit in the 10 day period when patient should isolate/mask, don full PPE (N-95 mask, eye protection/face shield, gloves, gown, and have patient don N-95 mask). Only the nurse and patient should be in the room during the visit.

**QUESTION 3B**

**If moderately / severely immunocompromised**

- 20 days from symptom start date: \_\_\_\_\_ (day 0; day 1 = the next full day)
- Advise patient to test as soon as possible.
- Advise patient to isolate from others and wear a well-fitting mask around others x 20 days since symptom onset (day 0 = symptom start date; day 1 = the next full day)
- Advise patient to avoid places where unable to wear a mask.
- End isolation/masking after 20 days if 1) At least 24 hours since last fever (without use of anti-pyretics) **and** 2) Symptoms have improved.
- May end isolation/masking earlier with testing if 1) At least 24 hours since last fever (without use of anti-pyretics) **and** 2) Symptoms have improved **and** 3) If using, a single negative Nucleic Acid Amplification Test (NAAT)/ Polymerase Chain Reaction (PCR) test is sufficient in most circumstances. If a higher level of clinical suspicion for COVID-19 infection exists, consider maintaining extending masking/isolation and confirming with a second negative NAAT/PCR test or if using an antigen test, a negative result should be confirmed by either a negative NAAT/PCR test or second negative antigen test taken 48 hours after the first negative test.
- Inform Nufactor
- If symptoms worsen or recur, restart isolation at day 0.
- If willing to make visit in the 20 day period when patient should isolate/mask, don full PPE (N-95 mask, eye protection/face shield, gloves, gown, and have patient don N-95 mask). Only the nurse and patient should be in the room during the visit.

Patient Name: \_\_\_\_\_

#### QUESTION 4

**Have you been in close contact with anyone who tested positive for COVID-19 in the past 10 days?** Close contact is being within 6 feet for a cumulative total of 15 minutes or more over a 24-hour period. YES (continue) NO (proceed to Part 2)

**Asymptomatic** (proceed)      **Symptomatic** (follow Question 2 guidance)

**Date of exposure:** \_\_\_\_\_

- 10 days from exposure date: \_\_\_\_\_ (day 0; day 1 = the next full day)
- Advise patient to isolate from others and wear a well-fitting mask around others x 10 days since exposure.
- Advise patient to avoid places where unable to wear a mask.
- If symptoms never develop, end isolation/masking after 10 days.
- If testing (for those who have not recovered from COVID-19 within past 30 days), may end masking earlier if:
  - Test (antigen test) immediately (but not earlier than 24 hours after the exposure). If negative, test (antigen test) again 48 hours after the first negative test. If negative, test (antigen test) again 48 hours after the second negative test. This will typically be at day 1 (where day of exposure is day 0), day 3, and day 5. If final test negative, can cease isolating/masking after day 7.
  - Testing generally not recommended if recovered from COVID-19 in prior 30 days.
  - Consider testing if recovered from COVID-19 in the prior 31-90 days; antigen test (instead of a Nucleic Acid Amplification Test [NAAT]/Polymerase Chain Reaction [PCR]) is recommended (some people may remain positive but not be infectious during this period).
- If symptoms develop, restart isolation at day 0.
- Inform Nufactor
- If willing to make visit in the 10 day period when patient should isolate/mask, don full PPE (N-95 mask, eye protection/face shield, gloves, gown, and have patient don N-95 mask). Only the nurse and patient should be in the room during the visit.

Patient Name: \_\_\_\_\_

**Part 2. Patient Household Member Screen**

Perform at same time as patient screen.  
Screen each household member.

**QUESTION 1**

Does any household member have any COVID-19 symptoms that are not caused by another condition such as allergies? Contact Nufactor if unsure.

**YES NO**

**If YES,**

- Only patient/nurse should be in the same room/area.

**QUESTION 2**

Has any household member been in close contact with anyone with COVID-19 in the past 10 days?

**YES NO**

Close contact is being within 6 feet for a cumulative total of 15 minutes or more over a 24-hour period with someone who has COVID-19.

**If YES,**

- Only patient/nurse should be in the same room/area.

**QUESTION 3**

Has any household member had a positive COVID-19 test for active virus in the past 20 days, or are they awaiting results of a COVID-19 test?

**YES NO**

**If YES,**

- Only patient/nurse should be in the same room/area
- Follow question 4 guidance.

**Part 3. Agency Nurse Self-Screen**

Perform before making a visit.

**QUESTION 1**

Do you have any COVID-19 symptoms that are not caused by another condition such as allergies? Contact Nufactor if unsure.

**YES NO**

**If yes, check all that apply:**

- Fever or chills
- Cough
- Shortness of breath or difficulty breathing
- Fatigue
- Muscle or body aches
- Headache
- New loss of taste or smell
- Sore throat
- Congestion or runny nose
- Nausea or vomiting
- Diarrhea

**If YES,**

- Inform Nufactor.
- Contact your manager to arrange restaffing of the visit.

**QUESTION 2**

Have you been in close contact with anyone with COVID-19 in the past 10 days?

**YES NO**

Close contact is being within 6 feet for a cumulative total of 15 minutes or more over a 24-hour period with someone who has COVID-19.

**If YES,**

- Do not proceed with the visit.
- Inform Nufactor.
- Contact your manager to arrange restaffing of the visit.

**QUESTION 3**

Have you had a positive COVID-19 test for active virus in the past 20 days, or are you awaiting results of a COVID-19 test?

**YES NO**

**If YES,**

- Do not proceed with the visit.
- Inform Nufactor.
- Contact your manager to arrange restaffing of the visit.

Reference: Infection Control: Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) CDC dated 03.18.24